



1. [21 points] Evaluate each limit.

$$(a) \lim_{x \rightarrow 7} \frac{x^2 - 8x + 7}{x^2 - 6x - 7} = \lim_{x \rightarrow 7} \frac{\cancel{(x-7)}(x-1)}{\cancel{(x-7)}(x+1)} = \frac{7-1}{7+1} = \frac{6}{8}$$

$$= \boxed{3/4}$$

$$(b) \lim_{x \rightarrow 1} \frac{\sqrt{4-3x} - 1}{x^2 - 1} = \lim_{x \rightarrow 1} \frac{(\sqrt{4-3x} - 1) \cdot (\sqrt{4-3x} + 1)}{(x+1)(x-1) \cdot (\sqrt{4-3x} + 1)}$$

$$= \lim_{x \rightarrow 1} \frac{(4-3x) - 1}{(x+1)(x-1)(\sqrt{4-3x} + 1)} = \frac{-3(x-1)}{(x+1)\cancel{(x-1)}(\sqrt{4-3x} + 1)}$$

$$= \frac{-3}{2 \cdot (\sqrt{1} + 1)} = \boxed{-3/4}$$

$$(c) \lim_{x \rightarrow 4^-} \frac{|x-4|}{x^2 - 3x - 4} \quad |x-4| = -(x-4) \text{ for } x < 4, \text{ so}$$

$$= \lim_{x \rightarrow 4^-} \frac{-(x-4)}{(x+1)(x-4)} = \frac{-1}{5}$$

$$= \boxed{-1/5}$$

(continued on reverse)

$$(d) \lim_{x \rightarrow -\infty} \frac{3 - 4x - 4x^2}{9 + 2x^2}$$

$$= \lim_{x \rightarrow -\infty} \frac{3 - 4x - 4x^2}{9 + 2x^2} \cdot \frac{1/x^2}{1/x^2}$$

$$= \lim_{x \rightarrow -\infty} \frac{3/x^2 - 4/x - 4}{9/x^2 + 2} = \frac{3/\infty - 4/(-\infty) - 4}{9/\infty + 2}$$

$$= \frac{-4}{2} = \boxed{-2}$$

$$(e) \lim_{x \rightarrow \infty} \frac{x^3 - 10x^2}{5x^2 + 7}$$

$$= \lim_{x \rightarrow \infty} \frac{x^3 - 10x^2}{5x^2 + 7} \cdot \frac{1/x^2}{1/x^2} = \lim_{x \rightarrow \infty} \frac{x - 10}{5 + 7/x^2}$$

$$= \frac{\infty - 10}{5 + 7/\infty} = \frac{\infty - 10}{5 + 0} = \boxed{+\infty}$$

$$(f) \lim_{x \rightarrow 8} \frac{\frac{x}{x+4} - \frac{x-4}{x-2}}{x^2 - 10x + 16}$$

$$= \lim_{x \rightarrow 8} \frac{\frac{x}{x+4} \cdot \frac{x-2}{x-2} - \frac{(x-4)}{x-2} \cdot \frac{x+4}{x+4}}{(x-2)(x-8)}$$

$$= \lim_{x \rightarrow 8} \frac{\frac{x^2 - 2x - x^2 + 16}{(x+4)(x-2)}}{(x-2)(x-8)}$$

$$= \lim_{x \rightarrow 8} \frac{-2(x-8)}{(x+4)(x-2)(x-2)(x-8)} = \frac{-2}{12 \cdot 6 \cdot 6}$$

$$= -\frac{1}{6 \cdot 6 \cdot 6} = \boxed{-\frac{1}{216}}$$

$$(g) \lim_{x \rightarrow 2^+} \frac{x^2 - x - 6}{x^2 + x - 6}$$

$$= \lim_{x \rightarrow 2^+} \frac{(x+2)(x-3)}{(x-2)(x+3)} = \frac{4 \cdot (-1)}{0^+ \cdot 5} = \frac{-4}{0^+} = \boxed{-\infty}$$

2. [15 points] Evaluate the derivative of each function. You do not need to simplify your answer.

(a)  $f(x) = (3x - 7) \left( x^{1/3} + \frac{1}{x^4} \right)$

$$f'(x) = 3 \cdot \left( x^{1/3} + \frac{1}{x^4} \right) + (3x - 7) \cdot \left( \frac{1}{3} x^{-2/3} - 4 \cdot \frac{1}{x^5} \right)$$

(b)  $g(x) = (2x^3 + 5x^4)^{1/3}$

$$g'(x) = \frac{1}{3} \cdot (2x^3 + 5x^4)^{-2/3} \cdot (6x^2 + 20x^3)$$

(c)  $h(x) = (x^2 + 7)\sqrt{5x + 3}$

$$h'(x) = 2x\sqrt{5x+3} + (x^2+7) \cdot \frac{1}{2\sqrt{5x+3}} \cdot 5$$

(continued on reverse)

$$(d) f(x) = \frac{\sqrt{2x+3}}{x^2+1}$$

$$f'(x) = \frac{\frac{1}{2\sqrt{2x+3}} \cdot 2 \cdot (x^2+1) - \sqrt{2x+3} \cdot 2x}{(x^2+1)^2}$$

$$(e) g(x) = (2x-1)^3(5x+3)^5$$

$$g'(x) = 3(2x-1)^2 \cdot 2 \cdot (5x+3)^5 + (2x-1)^3 \cdot 5(5x+3)^4 \cdot 5$$

3. [9 points] Let  $f(x) = \frac{2x}{3x+1}$ . Compute  $f'(x)$  using the limit definition of the derivative. You may use the quotient rule to check your answer, but for full points all steps of the limit calculation must be shown.

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{2(x+h)}{3(x+h)+1} - \frac{2x}{3x+1}}{h} = \lim_{h \rightarrow 0} \frac{\frac{2(x+h)(3x+1) - 2x[3(x+h)+1]}{[3(x+h)+1] \cdot (3x+1)}}{h} \\ &= \lim_{h \rightarrow 0} \frac{6x^2 + 2x + 6xh + 2h - 6x^2 - 6xh - 2x}{h(3(x+h)+1)(3x+1)} \\ &= \lim_{h \rightarrow 0} \frac{2h}{h(3(x+h)+1)(3x+1)} \\ &= \frac{2}{(3(x+h)+1)(3x+1)} \\ &= \boxed{\frac{2}{(3x+1)^2}} \end{aligned}$$

4. Consider the curve defined by the equation

$$y^2 = x^3 - x + 1.$$

- (a) [4 points] Determine  $\frac{dy}{dx}$  using **implicit differentiation**. Your answer will be in terms of both  $x$  and  $y$ .

$$\frac{d}{dx}(y^2) = \frac{d}{dx}(x^3 - x + 1)$$

$$2y \cdot \frac{dy}{dx} = 3x^2 - 1$$

$$\boxed{\frac{dy}{dx} = \frac{3x^2 - 1}{2y}}$$

- (b) [4 points] Find the equation of the tangent line at the point  $(3, 5)$ .

$$\text{@}(3, 5), \quad \frac{dy}{dx} = \frac{3 \cdot 3^2 - 1}{2 \cdot 5} = \frac{26}{10} = 13/5$$

So the tangent line is

$$y - 5 = \frac{13}{5}(x - 3)$$

ie.  $y = \frac{13}{5}x - \frac{39}{5} + 5$

ie.  $\boxed{y = \frac{13}{5}x - \frac{14}{5}}$

5. [9 points] Find the absolute maximum and absolute minimum values of  $f(x) = x^2(x-5)^3$  on the interval  $[0, 6]$ .

$$\begin{aligned}f'(x) &= 2x(x-5)^3 + x^2 \cdot 3(x-5)^2 \\&= x(x-5)^2 \cdot [2(x-5) + 3x] \\&= x(x-5)^2(5x-10) \\&= 5x(x-5)^2(x-2)\end{aligned}$$

$\Rightarrow$  crit. numbers  $0, 2, 5$ .

Closed interval method:

$$f(0) = 0^2 \cdot (-5)^3 = 0$$

$$f(2) = 2^2 \cdot (-3)^3 = 4 \cdot (-27) = -108 \quad \leftarrow \text{min}$$

$$f(5) = 5^2 \cdot 0^3 = 0$$

$$f(6) = 6^2 \cdot 1^3 = 36 \quad \leftarrow \text{max}$$

abs. max. is 36 @  $x=6$   
abs. min. is -108 @  $x=2$ .

6. [12 points] Consider the following function.

$$f(x) = \frac{2-x}{x^2}$$

inc/dec/intro	4
conc./inf.	4
amp.	4

Sketch the graph  $y = f(x)$ . Clearly label the following features on your graph: asymptotes (horizontal or vertical), intervals where it is increasing/decreasing, intervals where it is concave up/down, local max(s) and min(s), and inflection point(s).

Division by 0 occurs @  $x=0$ , & numerator is  $2-0 \neq 0$  then  
 $\Rightarrow$  V.A. @  $x=0$ .

$$\lim_{x \rightarrow \infty} \frac{2-x}{x^2} = \lim_{x \rightarrow \infty} \left( \frac{2}{x^2} - \frac{1}{x} \right) = \frac{2}{\infty} - \frac{1}{\infty} = 0 \quad \left. \vphantom{\lim_{x \rightarrow \infty} \frac{2-x}{x^2}} \right\} \text{so H.A. @ } y=0$$

$$\& \lim_{x \rightarrow -\infty} \frac{2-x}{x^2} = \frac{2}{(-\infty)} - \frac{1}{-\infty} = 0$$

$$f'(x) = \left( \frac{2}{x^2} - \frac{1}{x} \right)' = -\frac{4}{x^3} - (-1)\frac{1}{x^2} = -\frac{4}{x^3} + \frac{1}{x^2} = \frac{x-4}{x^3} \quad \left. \vphantom{f'(x)} \right\} \begin{array}{l} \text{num. 0 @ } x=4 \\ \text{denom 0 @ } x=0 \end{array}$$

	$\longleftarrow \begin{array}{c} 0 \qquad 4 \\ \hline \end{array} \longrightarrow$		
$x-4$	-	-	+
$\frac{1}{x^2}$	-	+	+
$f'$	+	-	+
$f$	↗	↘	↗

$\Rightarrow$   $\left\{ \begin{array}{l} \text{inc. on } (-\infty, 0) \text{ \& } (4, \infty) \\ \text{dec. on } (0, 4) \\ \text{Local min @ } x=4, y=f(4) = -1/8 \\ \text{no extremum @ } x=0 \text{ (it's a discontinuity).} \end{array} \right.$

$$f''(x) = \left( -\frac{4}{x^3} + \frac{1}{x^2} \right)' = \frac{12}{x^4} - \frac{2}{x^3} = \frac{12-2x}{x^4} = -2 \cdot \frac{x-6}{x^4} \quad \left. \vphantom{f''(x)} \right\} \begin{array}{l} \text{num. 0 @ } x=6 \\ \text{denom 0 @ } x=0 \end{array}$$


	$\longleftarrow \begin{array}{c} 0 \qquad 6 \\ \hline \end{array} \longrightarrow$		
$x-6$	-	-	+
$\frac{1}{x^4}$	-	-	-
$f''$	+	+	-
$f$	∪	∪	∩

$\Rightarrow$   $\left\{ \begin{array}{l} \text{conc. up on } (-\infty, 0) \text{ \& } (0, 6) \\ \text{conc. down on } (6, \infty) \\ \text{inflection pt. @ } x=6, y=f(6) = -\frac{4}{36} = -1/9. \end{array} \right.$

(sketch on next page)

This page intentionally left blank. You may use it for scratchwork.

Sketch: (not exactly to scale)




7. [12 points] A 15 foot ladder is leaning against a wall, and sliding down the side. At this moment, the bottom of the ladder is 9 feet away from the wall, and is sliding away at 2 feet per second. Determine how quickly the top of the ladder is sliding down the wall at this moment (in feet per second).



$$x^2 + y^2 = 15^2$$

At key moment



$$x = 9 \quad y = \sqrt{15^2 - 9^2} = 12$$

$$x' = 2 \quad \text{want: } y' = ?$$

$$\frac{d}{dt}(x^2 + y^2) = \frac{d}{dt}(15^2)$$

$$\Rightarrow 2x \cdot x' + 2y \cdot y' = 0$$

$\Rightarrow$  At key moment,

$$2 \cdot 9 \cdot 2 + 2 \cdot 12 \cdot y' = 0$$

$$24y' = -36$$


$$y' = -\frac{36}{24} = \underline{-3/2}$$

(neg. since decreasing as ladder falls).

The top of the ladder is falling  
at 3/2 foot per second.

8. [12 points] A small rectangular box with a square base **with no lid** is to be constructed out of 12 square inches of cardboard. Determine what dimensions the box should have in order for its volume to be as large as possible.

Note: this is problem 3.7.15, which was on the homework (with smaller numbers to make the arithmetic easier).



cardboard needed:

$$\underbrace{x^2}_{\text{base}} + \underbrace{4xy}_{\text{4 sides of area } xy} = 12 \quad (\text{constraint})$$

We want to maximize volume:

$$V = x^2 y.$$

Solve for  $y$  using the constraint:

$$4xy = 12 - x^2$$

$$y = \frac{12 - x^2}{4x}$$

So Volume, as a function of  $x$ , is  $V(x) = x^2 \left( \frac{12 - x^2}{4x} \right) = \frac{1}{4} x (12 - x^2)$

Feasible values: we need  $x > 0$

$$\& y > 0 \Rightarrow \frac{12 - x^2}{4x} > 0 \Rightarrow 12 > x^2 \Rightarrow x < \sqrt{12} = 2\sqrt{3}$$

$\Rightarrow$  the interval is  $[0, 2\sqrt{3}]$ .

We can maximize w/ the closed interval method:

$$V(x) = \frac{1}{4} x (12 - x^2)$$

$$V'(x) = \frac{1}{4} \cdot 1 \cdot (12 - x^2) + \frac{1}{4} x \cdot (-2x) = \frac{1}{4} (12 - x^2 - 2x^2)$$

$$= \frac{1}{4} (12 - 3x^2) = \frac{3}{4} (4 - x^2)$$

$\Rightarrow$  crit. pts @  $x = \pm 2$ ; discard  $-2$  since it's out of the interval.


Check endpoints &  $x=2$ :

$$V(0) = \frac{1}{4} \cdot 0 \cdot 12 = 0$$

$$V(2) = \frac{1}{4} \cdot 2 \cdot 8 = 4 \leftarrow \underline{\underline{\text{max}}}$$

$$V(2\sqrt{3}) = \frac{1}{4} \cdot 2\sqrt{3} \cdot 0 = 0$$


So the max volume is 4, which occurs for  $x=2$ ,  $y = \frac{12 - 2^2}{4 \cdot 2} = 1$ .



9. Consider the following piecewise function.

$$f(x) = \begin{cases} 2 & x < 0 \\ \frac{1}{x-2} & 0 \leq x < 2 \\ 2\sqrt{x-2} & x \geq 2 \end{cases}$$

- (a) [6 points] Sketch the graph  $y = f(x)$  on the axes below (for this sketch, you don't need to take any derivatives or apply the techniques from Chapter 3; instead think about how each piece of the graph is obtained from a graph you already know about).



- (b) [6 points] Evaluate each of the following quantities (no explanation or scratchwork is required).

$$\bullet \lim_{x \rightarrow 11^-} f(x) = 2$$

$$\bullet \lim_{x \rightarrow 2^-} f(x) = -\infty$$

$$\bullet \lim_{x \rightarrow 0} f(x) = -\frac{1}{2}$$

$$\bullet \lim_{x \rightarrow 2^+} f(x) = 0$$

$$\bullet f(0) = -\frac{1}{2}$$

$$\bullet f(2) = 0$$

(continued on reverse)

The definition of  $f(x)$  is reproduced below for convenience.

$$f(x) = \begin{cases} 2 & x < 0 \\ \frac{1}{x-2} & 0 \leq x < 2 \\ 2\sqrt{x-2} & x \geq 2 \end{cases}$$

(c) [3 points] Determine all points where  $f(x)$  is discontinuous.

$x=0$  is a jump discontinuity  
( $\lim_{x \rightarrow 0^-} f(x) \neq \lim_{x \rightarrow 0^+} f(x)$ )

$x=2$  is an infinite discontinuity  
( $\lim_{x \rightarrow 2^-} f(x) = -\infty$ )

There are no other discontinuities;  
the three functions used are continuous  
on the intervals where they apply.

(continued on reverse)

10. [12 points] Consider the function

$$f(x) = \frac{x}{4+x^2}$$

(a) Compute  $f'(x)$ , and simplify.

$$f'(x) = \frac{1 \cdot (4+x^2) - x \cdot 2x}{(4+x^2)^2}$$

either of these is a fine way to leave the answer.

$$\left\{ \begin{aligned} &= \frac{4-x^2}{(4+x^2)^2} \\ &= -\frac{(x+2)(x-2)}{(4+x^2)^2} \end{aligned} \right.$$

(b) Determine all critical numbers of  $f(x)$ .

denom. <sup>of  $f'(x)$</sup>  is never 0 since  $4+x^2 > 0$ .  
 num. <sub>of  $f'(x)$</sub>  is 0 @  $\boxed{x = \pm 2}$ .

(continued on reverse)

(c) Determine the intervals on which  $f(x)$  is increasing and decreasing.

	$\xleftarrow{-2} \quad \quad \quad \xrightarrow{2}$		
$-\frac{1}{(4+x^2)^2}$	-	-	-
$x+2$	-	+	+
$x-2$	-	-	+
$f'$	-	+	-
$f$	$\searrow$	$\nearrow$	$\searrow$

dec. on  $(-\infty, -2)$  &  $(2, \infty)$   
 inc. on  $(-2, 2)$

(d) Classify each critical point as a local minimum, a local maximum, or neither.

$x = -2$  is a local min.  
 by 1<sup>st</sup> deriv. test  $(\searrow \nearrow)$

$x = +2$  is a local max  
 by 1<sup>st</sup> deriv. test  $(\nearrow \searrow)$ .